

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 186 294 A2**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
13.03.2002 Bulletin 2002/11

(51) Int Cl.7: **A61K 31/215, A61K 31/19**

(21) Application number: **01118978.4**

(22) Date of filing: **06.08.2001**

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR**
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: **07.08.2000 JP 2000238039**

(71) Applicant: **Kao Corporation**
Tokyo 103-8210 (JP)

(72) Inventors:
• **Suzuki, Atsushi**
Haga-gun, Tochigi 321-3497 (JP)
• **Ochiai, Ryuji**
Haga-gun, Tochigi 321-3497 (JP)
• **Tokimitsu, Ichiro**
Haga-gun, Tochigi 321-3497 (JP)

(74) Representative: **VOSSIUS & PARTNER**
Siebertstrasse 4
81675 München (DE)

(54) **Ferulic acid and caffeic or chlorogenic acids containing compositions for alleviating hypertension or preventing a rise in blood pressure**

(57) Products and compositions for preventing or reducing the severity of hypertension. These products contain (a) ferulic acid or a ferulate ester, and (b) caffeic acid and/or a chlorogenic acid. The preventive or rem-

edy can suppress a rise in blood pressure and alleviate hypertension, and is usable as a food.

EP 1 186 294 A2

Description

[0001] The present invention relates to products and compositions that prevent, remedy or reduce the severity of hypertension and that are capable of suppressing a rise in blood pressure.

[0002] Hypertension in Japan ranks first among reasons why patients attend hospitals. According to the National Life Fundamental Survey of Ministry of Health and Welfare (fiscal 1998), in Japan, 64 patients per 1000 were admitted to hospitals for hypertension.

[0003] Heart diseases such as angina pectoris, myocardial infarction and heart failure and cerebrovascular diseases such as cerebral infarction, cerebral hemorrhage and subarachnoid hemorrhage are closely related to hypertension and rank second and third, respectively, among the causes of death of the Japanese.

[0004] Hypertension may be treated by the administration of blood-pressure lowering pharmaceuticals such as diuretics, sympathetic inhibitors, vasodilators or angiotensin-converting enzyme inhibitors. Such drugs are mainly applied to patients suffering from severe hypertension. Although many of the pharmaceuticals administered to treat hypertension are satisfactory in their effectiveness, significant side-effects such as tachycardia and bradycardia can be a serious burden for patients.

[0005] Hypertension, especially its milder forms, may also be treated by generally improving lifestyle, such as through dietetic therapy, kinesitherapy and limitation of alcoholic intake or smoking. The importance of such changes in lifestyle is now being increasingly recognized and appreciated, not only for milder forms of hypertension, but also for more severe cases.

[0006] Above all, improvement of eating habits has received great attention. There exist a large number of foods, which have traditionally been said to have blood pressure lowering action. Food products have been briskly searched in order to identify and isolate components that lower blood pressure.

[0007] It has been reported that phenols such as caffeic acid contained crude form in the spike of *Schizonepeta tenuifolia* Briq. exerts calcium antagonism and may be useful for the treatment of vascular diseases such as hypertension (Japanese Patent Application Laid-Open (*Kokai*) No. Hei 4-243822).


[0008] The use the juice of an unripe fruit of apples, pears, peaches or the like fruits belonging to the family *Rosaceae* has also been proposed a hypotensor. Such juice contains, as polyphenols, caffeic acid and chlorogenic acid having angiotensin I converting enzyme (ACE) inhibitory action (Japanese Patent Application Laid-Open (*Kokai*) No. Hei 8-219453).



[0009] However, foods which are said to be effective for lowering blood pressure or their effective ingredients are not always satisfactory in their effectiveness and many of them do not start to exert significant blood pressure reducing effects immediately after intake or exert long-last anti-hypertensive effects.

[0010] Therefore, one object of the present invention is to provide a preventive or remedy for hypertension which has excellent safety, does not become a burden for patients even by daily intake, has higher antihypertensive action and exerts significant prompt and/or long-lasting antihypertensive effects.

[0011] The present inventors have found that a combination of ferulic acid (or an ester thereof), and caffeic acid and/or chlorogenic acid exerts prompt blood pressure lowering effects and, by prolonged administration, suppresses rises in blood pressure. This combination of ingredients also has reduced side effects such as bradycardia.

[0012] The chemical structures of caffeic acid, one type of a chlorogenic acid, and ferulic acid are shown below:

Caffeic acid

Chlorogenic acidFerulic acid

[0013] The present invention thus provides products and compositions for the prevention, alleviation or reduction of hypertension. These compositions comprise the following components (a) and (b):

- (a) a component selected from ferulic acid, an ester thereof or a pharmaceutically acceptable salt thereof, and
- (b) a component selected from caffeic acid or a chlorogenic acid, or a pharmaceutically acceptable salt thereof.

[0014] Another aspect of the present invention, provides a food containing or supplemented to contain the above-described components (a) and (b).

[0015] A further aspect of the present invention provides the use of the above-described components (a) and (b) for the preparation of a product that prevents, treats, reduces or remedies hypertension.

[0016] A still further aspect of the present invention provides a method for treating hypertension that comprises the administration of an effective amount of the above-described components (a) and (b).

[0017] A further aspect of the invention is a method for providing prompt blood pressure reduction by the administration of a composition comprising ferulic acid, an ester thereof, or a pharmaceutically acceptable salt thereof.

[0018] A still further aspect of the invention is a method for providing long-lasting blood pressure reduction by administering a composition comprising caffeic acid or a chlorogenic acid.

[0019] The ferulic acid or ester thereof as component (a) to be used in the present invention can be obtained by either extraction from a natural substance, particularly, a plant which contains it, or by its industrial preparation, for instance, by chemical synthesis.

[0020] Preferred examples of plants containing ferulic acid or esters of ferulic acid include coffee, onion, Japanese radish, lemon, *Angelicae radix*, *Cnidii Rhizoma*, goldthread, asafoetida, sugarcane, corn, barley and rice, with rice being particularly preferred. The term "rice" as used herein means raw or dry seeds of rice (*Oryza sativa* LINNE).

[0021] Esters of ferulic acid include those obtained by conversion upon extraction or fractionation of those originally contained in a natural substance, particularly, a plant; and to the chemically modified products thereof. For example, rice bran oil obtained from rice bran is separated using hydrous ethanol and hexane at room temperature under a weak alkaline condition and ferulate ester is available in the hydrous ethanol fraction. Ferulic acid can be obtained by hot hydrolysis of the ferulate ester obtained in the above-described manner with sulfuric acid under pressure, followed by purification. It can also be obtained by culturing bacteria (*Pseudomonas*) in a broth containing a clove oil obtained by steam distillation of buds and leaves of *Syzygium aromaticum* MERRILL et PERRY or a broth containing eugenol

available by purification of the clove oil, followed by separation of the resulting culture broth and purification. Chemical synthesis of ferulic acid is attained, for example, by condensation reaction of vanillin and malonic acid, Journal of American Chemical Society, 74: 5346 (1952). Ferulic acid has steric isomers. Any one of them is usable. A mixture of the isomers is also usable.

[0022] As the alcohol moiety of the ferulate ester to be used in the present invention, C₁₋₄₀ alcohols are preferred. Examples include linear or branched alkyl or alkenyl alcohols, aryl alcohols, monoterpene alcohols, sesquiterpene alcohols, diterpene alcohols, triterpene alcohols, sterols, and trimethyl sterols, more specifically, ethanol, oleyl alcohol, 2-ethyl-hexyl alcohol, allyl alcohol, cetyl alcohol, menthyl alcohol, phenol, benzyl alcohol, cholesterol, cycloartenol, 24-methylene cycloartenol, campesterol, β -sitosterol, cycloartanol, cycloprenol, α -sitosterol, stigmasterol, stigmastanol, α -sitostanol, β -sitostanol and campestanol.

[0023] Ferulic acid has improved water solubility and increased physiological availability when it is in the form of a salt. No particular limitation is imposed on the salt of the ferulic acid insofar as it is pharmaceutically acceptable. Examples of a basic substance for the formation of such a salt include hydroxides of an alkali metal such as lithium hydroxide, sodium hydroxide and potassium hydroxide, hydroxides of an alkaline earth metal such as magnesium hydroxide and calcium hydroxide, inorganic bases such as ammonium hydroxide and basic amino acids such as arginine, lysine, histidine and ornithine, and organic bases such as monoethanolamine, diethanolamine and triethanolamine. Out of them, hydroxides of an alkali metal or alkaline earth metal are particularly preferred. As a preventive or remedy for hypertension according to the present invention, such a salt, which has been prepared in advance, may be added to a composition composed of the other components, or ferulic acid and a salt-forming component therewith may be added to the composition separately to form their salt therein.

[0024] As Component (a), at least two forms of ferulic acid or a ferulic acid derivative or salt may be used in combination. It is preferred for an adult (weight: 60 kg) to take Component (a) in an amount of about 0.001 to about 10 g, preferably about 0.005 to about 5 g, more preferably about 0.01 to about 0.5 g a day.

[0025] The caffeic acid or chlorogenic acid to be used as Component (b) in the present invention can be obtained by extraction from a natural substance, particularly, a plant containing these substances, or by industrial preparation, for instance, by chemical synthesis. Examples of the chlorogenic acid to be used in the present invention include neochlorogenic acid, isochlorogenic acids (such as 3,5-dicaffeoylquinic acid), cryptochlorogenic acid, feruloylquinic acid and 5-caffeoylquinic acid. Component (b) may be an extract of a plant abundant in chlorogenic acid such as raw coffee beans, leaves of a nandina and unripe apple fruits, or alternatively, an extract of raw coffee beans available by extraction of the seeds of *Coffea arabica* LINNE with an aqueous solution of ascorbic acid or citric acid under warming.

[0026] The caffeic acid and chlorogenic acid products forming Component (b), like those of ferulic acid, have improved water solubility and increased physiological availability when they are in the form of a pharmaceutically acceptable salt. As the salt as component (b), those exemplified above as the salts of ferulic acid can be mentioned. It is preferred for an adult (weight: 60 kg) to take Component (b) in an amount of about 0.001 to about 10 g, more preferably about 0.005 to about 5 g a day.

[0027] As Component (b), at least two derivatives or salts of a chlorogenic acid or a caffeic acid product may be used in combination.

[0028] The preventive or remedy for hypertension according to the present invention can be formed into an orally administrable or parenterally administrable composition by adding to its effective ingredient a pharmaceutically acceptable carrier. Examples of the orally administrable composition include tablets, granules, fine subitlaes, pills, powders, capsules (hard capsules and soft capsules), troches, chewables and liquids (medical drinks). Examples of the parenterally administrable composition include intravenously administrable preparations such as isotonic, sterile solutions for injection, suppositories and dermatologic preparations for external use.

[0029] The compositions for preventing or treating hypertension or high blood pressure according to the present invention have a high degree of safety so that no problem occurs even if those who have a normal blood pressure usually take it as a food. Examples of such compositions in the form of food or beverage include beverages such as juice and coffee, liquid foods such as soup, emulsion or pasty foods such as milk and curry, semisolid foods such as jelly and gummy; solid foods such as gum, tofu and supplement; powdery foods; and oil or fat containing foods such as margarine, mayonaise and dressing.

[0030] The weight ratio of Component (a) to Component (B) in a combined preparation, that is, Component (a)/(b) preferably ranges from about 0.01 to 50, more preferably ranges from about 0.01 to 5. The weight ratio of Component (a) to Component (b) is confirmed by high-performance liquid chromatography equipped with an electrochemical detector.

[0031] It is preferred for an adult (weight: 60 kg) to take the preventive or remedy for hypertension according to the present invention so that the total amount of Components (a) and (b), the effective ingredients, would be about 0.001 to about 20 g, particularly about 0.005 to about 10 g a day. When a plant extract is employed, it is preferred to take it in an amount in terms of a dry weight.

[0032] In addition to the combined preparations of the present invention comprising ferulic acid and caffeic acid and/

or a chlorogenic acid, compositions comprising any of these ingredients may be formulated to decrease the effects of hypertension or reduce high blood pressure.

[0033] Foods or beverages associated with hypertension may advantageously be supplemented with caffeic acid, a chlorogenic acid and/or ferulic acid in dosages that preferably inhibit or reduce the hypertensive effects of the food or beverage. For instance, beverages containing caffeine, such as coffee, have been associated with hypertension and may be supplemented with amounts of caffeic acid, chlorogenic acid and/or ferulic acid to reduce hypertensive effects associated with the consumption of these beverages.

[0034] Caffeic acid, a chlorogenic acid and/or ferulic acid may also be compounded as food or nutritional supplements in amounts which preferably reduce hypertension. For instance, these substances may be admixed with a pharmaceutically acceptable excipient, filler or carrier. As such, they may be placed in tablet or capsule form, or compounded in standardized dosages in solid or liquid form.

[0035] Breeding or engineering of agricultural products to contain or express higher amounts of caffeic acid, chlorogenic acid and/or ferulic acid may be done so as to obtain products that exert beneficial effects in hypertensive subjects.

Examples

Example 1: Evaluation of blood pressure reduction

(1) Animals provided for test

[0036] Prior to initiation of the experimental studies in order to accustom the test animals to sphygmomanometric operation the blood pressure of each test animal--male, 15 week-old, spontaneously hypertensive rats ("SHR")--was preliminarily measured for 7 successive days using a commercially available noninvasive sphygmomanometer for rats (manufactured by Softlon Co., Ltd.).

[0037] Rats were all bred in a breeding room in a rat region under uniform conditions at a room temperature at $25 \pm 1^\circ\text{C}$, humidity of $55 \pm 10\%$ RH and illumination for 12 hours (from 7:00 am to 7:00 pm).

(b) Administration method and amount

[0038] In the control group (Control Plot, see Table 1), physiological saline was employed. As a material to be administered, a solution obtained by dissolving 0.2 wt.% (which will hereinafter be simply described as %) of caffeic acid in physiological saline, a solution obtained by dissolving 0.2 % of a chlorogenic acid in physiological saline, a solution obtained by dissolving 0.2 % of ferulic acid in physiological saline and a solution obtained by dissolving 0.1% of caffeic acid and 0.1% of ferulic acid in physiological saline were used in Test plot 1, Test plot 2, Test plot 3 and Test plot 4, respectively. In Test plot 5, a solution obtained by dissolving 0.1% of a chlorogenic acid and 0.1% of ferulic acid in physiological saline was used, while in Test plot 6, a solution obtained by dissolving 0.05% of caffeic acid, 0.05% of a chlorogenic acid and 0.1% of ferulic acid in physiological saline was used. Each of them was administered from the common carotid artery. The dosage was 1 mL/kg.

(c) Test method

[0039] SHRs were fasted overnight and employed for each test group, each group consisted of 5 rats. Systolic blood pressures of the caudal artery prior to intravenous administration, and 10 minutes and 1 hour each after administration were measured.

(4) Statistical treatment method

[0040] In Table 1, the changing ratio (%) in systolic blood pressure for each experimental group is reported by its mean value (%) and by its standard deviation. The Student's t-test was used to indicate the statistical significance of the experimental data. A significance level of 5% or less is indicated as "*" in Table 1, and significance level of 0.1% or less as "****".

[0041] A lowering ratio of the systolic blood pressure of each of 10 minutes and 1 hour after administration to the systolic blood pressure prior to administration is shown in Table 1. As is apparent from Table 1, the compositions of the present invention induced prompt and/or long-lasting antihypertensive effects and reductions in blood pressure.

Table 1

	Systolic blood pressure (changing ratio %)	
	after 10 minutes	After 1 hour
Control plot	saline	-1.9 \pm 1.4
Test plot 1	Caffeic acid (CA)	-10.2 \pm 0.5 ***
Test plot 2	Chlorogenic acid (CHA)	-7.2 \pm 1.7 *
Test plot 3	Ferulic acid (FA)	0.7 \pm 2.6
Test plot 4	CA + FA	-11.3 \pm 1.3 ***
Test plot 5	CHA + FA	-10.9 \pm 0.8 ***
Test plot 6	CA + CHA + FA	-13.6 \pm 3.4 ***

*, ***, A significance level was 5% or less and 0.1% or less, respectively relative to the control group, meaning existence of a significant difference.

EP 1 186 294 A2

[0042] Indicated is a mean value + standard deviation Example 2-Inhibition of Blood Pressure Rise

(1) Test Animals

5 [0043] Under similar conditions to Example 1, male, 6 week old spontaneously hypertensive rats (SHR) were bred.

(2) Administration method and amount

10 [0044] In the Control Plot, test animals were maintained on commercially available powdery feed and drinking water *ad libitum*.

[0045] In Test plot 1, they were maintained *ad libitum* on drinking water having 0.2% of sodium caffeinate added thereto and commercially available powdery feed having, incorporated therein, 1% of a triterpenol-ferulate ester mixture extracted from rice bran;

15 [0046] In Test Plot 2, on drinking water having 0.2% of a chlorogenic acid and commercially available powdery feed having, incorporated therein, 1% of a ferulic acid and cycloartenol ferulate;

[0047] In Test Plot 3, on drinking water having, added thereto, 0.2% of caffeic acid, 1% of a chlorogenic acid and 0.04% of ferulic acid.

(3) Test method

20 [0048] SHRs were employed for the test in groups, each group consisting of 6 rats. Systolic pressure of the caudal artery 4 weeks after administration was measured and the results were treated statistically in a similar manner to Example 1.

25 [0049] Systolic blood pressures prior to administration and 4 weeks after administration are shown in Table 2. Table 2 shows that the according to the present invention exert a marked inhibitory action against blood pressure rise.

30

35

40

45

50

55

Table 2

	Systolic blood pressure (mmHg)		
		Prior to administration	4 weeks after administration
Control plot	Water	152.1 \pm 4.4	201.0 \pm 3.9
Test plot 1	Na Caffeinate, triterpenol ferulate ester	155.3 \pm 3.7	183.5 \pm 4.0 *
Test plot 2	Chlorogenic acid, ferulic acid, cycloartenol ferulate	153.0 \pm 5.2	181.9 \pm 5.3 *
Test plot 3	Caffeic acid, chlorogenic acid, ferulic acid	155.0 \pm 4.1	188.2 \pm 3.4 *

*: The significance level relative to the control plot was 5% or less, indicating a significant lowering of the systolic blood pressure of groups treated with the compositions of the present invention.

Indicated is each mean \pm standard deviation.

Example 3: Soft capsules

[0050]

Gelatin	70.0%
Glycerin	22.9
Methyl paraoxybenzoate	0.15
Propyl paraoxybenzoate	0.51
Water	6.44

[0051] Soft, oval-type capsules having a weight of 150 mg and composed of the above-described composition were filled in a manner known *per se* in the art with 400 mg of soybean oil, 50 mg of caffeic acid and 50 mg of ferulic acid. These capsules when administered to test subjects exhibit good blood pressure lowering action.

Example 4- Beverage

[0052] A beverage according to the present invention was prepared by combining the ingredients enumerated below.

Skim milk	3.5%
Enzyme-hydrolyzed milk casein	3.5
Fructose	9.0
Chlorogenic acid	0.1
Sodium ferulate	10.0
Citric acid	0.1
Ascorbic acid	0.1
Flavor	0.1
Water	73.6

[0053] It has been found that the beverage having the above-described composition had high storage stability and had good taste.

Example 5 - Wheat Flour Products

[0054] Wheat flour products according to the present invention were prepared by the combination of the following ingredients:

Rapeseed oil	15 g
Corn starch	15
Wheat flour	42.6
Butter	5
Fructose	14
Ferulic acid	2
Cycloartenol ferulate	0.4
Table salt	0.5
Sodium bicarbonate	0.5
Water	5

[0055] Cookies having the above-described composition were baked.

Modifications and other embodiments

[0056] Various modifications and variations of the described anti-hypertensive products, compositions and methods, as well as the concept of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments,

it should be understood that the invention as claimed is not intended to be limited to such specific embodiments. Various modifications of the described modes for carrying out the invention which are obvious to those skilled in the medical, biological, chemical or pharmacological arts or related fields are intended to be within the scope of the following claims.

Incorporation by Reference

[0057] Each document, patent application or patent publication cited by or referred to in this disclosure is incorporated by reference in its entirety. Any patent document to which this application claims priority is also incorporated by reference in its entirety. Specifically, priority document JP, 2000-238039, filed August 7, 2000 is hereby incorporated by reference.

Claims

1. A composition comprising
 - (a) ferulic acid or an ester thereof, or a pharmaceutically acceptable salt thereof, and
 - (b) caffeic acid and/or a chlorogenic acid, or a pharmaceutically acceptable salt thereof.
2. The composition of claim 1, wherein (b) consists of caffeic acid or a pharmaceutically acceptable salt thereof.
3. The composition of claim 1, wherein (b) consists of a chlorogenic acid, or a pharmaceutically acceptable salt thereof.
4. The composition of claim 3, wherein the chlorogenic acid is selected from the group consisting of: 3,5-dicaffeoylquinic acid, cryptochlorogenic acid and 5-caffeoylquinic acid.
5. The composition of claim 1, wherein (b) comprises caffeic acid and a chlorogenic acid, or pharmaceutically acceptable salts thereof.
6. A food or beverage comprising a composition according to claim 1.
7. The food or beverage of claim 6, selected from the group consisting of a milk product, a cereal or pasta product, a soy product, a rice product, a wheat flour product, an oil or fat-containing food, a condiment, a semisolid food, a sauce, a soup, coffee, decaffeinated coffee, tea, decaffeinated tea, a herbal tea or infusion, a chocolate product, a beverage containing alcohol, a fruit or vegetable juice and a soft drink.
8. A supplement which is composition according to claim 1.
9. The supplement of claim 8, further comprising a nutrient, vitamin, mineral, herbal product or drug.
10. The supplement of claim 8 in the form of a tablet, granule, fine subtile, pill, powder, capsule, troche, medicinal drink, solution for injection, suppository, or dermatological preparation.
11. Use of
 - (a) ferulic acid or an ester thereof, or a pharmaceutically acceptable salt thereof, and
 - (b) a component selected from caffeic acid and/or a chlorogenic acid, or a pharmaceutically acceptable salt thereoffor the preparation of a pharmaceutical composition for preventing or treating hypertension or high blood pressure.
12. Use of claim 11, for reduction of systolic blood pressure.
13. Use of claim 11, for reduction of diastolic blood pressure.
14. Use of

EP 1 186 294 A2

(a) ferulic acid or an ester thereof, or a pharmaceutically acceptable salt thereof, and
(b) a component selected from caffeic acid and/or a chlorogenic acid, or a pharmaceutically acceptable salt thereof
for the preparation of a pharmaceutical composition for reducing a rise in blood pressure.

- 5
15. Use of claim 14, for reduction of the rise of systolic blood pressure.
16. Use of claim 14, for reduction of the rise in diastolic blood pressure.
- 10 17. A food or beverage supplemented with ferulic acid, or a pharmaceutically acceptable salt thereof, caffeic acid or a pharmaceutically acceptable salt thereof or with a chlorogenic acid, or a pharmaceutically acceptable salt thereof.

15

20

25

30

35

40

45

50

55

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 186 294 A3

(12)

EUROPEAN PATENT APPLICATION

(88) Date of publication A3:
21.05.2003 Bulletin 2003/21

(51) Int Cl.7: **A61K 31/215, A61K 31/19,
A61P 9/12**

(43) Date of publication A2:
13.03.2002 Bulletin 2002/11

(21) Application number: **01118978.4**

(22) Date of filing: **06.08.2001**

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR**
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: **07.08.2000 JP 2000238039**

(71) Applicant: **Kao Corporation**
Tokyo 103-8210 (JP)

(72) Inventors:
• **Suzuki, Atsushi**
Haga-gun, Tochigi 321-3497 (JP)
• **Ochiai, Ryuji**
Haga-gun, Tochigi 321-3497 (JP)
• **Tokimitsu, Ichiro**
Haga-gun, Tochigi 321-3497 (JP)

(74) Representative: **VOSSIUS & PARTNER**
Siebertstrasse 4
81675 München (DE)

(54) **Ferulic acid and caffeic or chlorogenic acids containing compositions for alleviating hypertension or preventing a rise in blood pressure**

(57) Products and compositions for preventing or reducing the severity of hypertension. These products contain (a) ferulic acid or a ferulate ester, and (b) caffeic acid and/or a chlorogenic acid. The preventive or rem-

edy can suppress a rise in blood pressure and alleviate hypertension, and is usable as a food.

EP 1 186 294 A3



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number

which under Rule 45 of the European Patent Convention shall be considered, for the purposes of subsequent proceedings, as the European search report

EP 01 11 8978

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (InCl.7)
X	ABRAHAM S K: "ANTI-GENOTOXIC EFFECTS IN MICE AFTER THE INTERACTION BETWEEN COFFEE AND DIETARY CONSTITUENTS" FOOD AND CHEMICAL TOXICOLOGY, XX, XX, vol. 34, no. 1, 1996, pages 15-20, XP001148404 ISSN: 0278-6915 * table 1 * * page 17, paragraph 1 *	1-10, 17	A61K31/215 A61K31/19 A61P9/12
X	US 4 707 472 A (AOKI HIDEMI ET AL) 17 November 1987 (1987-11-17)	17	
Y	* column 1, line 18 - line 23 *	1-16	
X	WO 98 01143 A (AAR PHARMA ; RUEPP MICHEL O (DE)) 15 January 1998 (1998-01-15)	17	
Y	* page 2, paragraph 1 * * page 11, paragraph 2 *	1-16	
X	US 5 958 417 A (HSU CHAU-SHIN) 28 September 1999 (1999-09-28)	17	
Y	* column 2, line 55 - line 63 *	1-16	
	-/--		
			TECHNICAL FIELDS SEARCHED (InCl.7)
			A61K
INCOMPLETE SEARCH			
<p>The Search Division considers that the present application, or one or more of its claims, does/do not comply with the EPC to such an extent that a meaningful search into the state of the art cannot be carried out, or can only be carried out partially, for these claims.</p> <p>Claims searched completely:</p> <p>Claims searched incompletely:</p> <p>Claims not searched:</p> <p>Reason for the limitation of the search:</p> <p>see sheet C</p>			
Place of search		Date of completion of the search	Examiner
THE HAGUE		26 March 2003	Leherte, C
CATEGORY OF CITED DOCUMENTS			
<p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons</p> <p>& : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03 02 (P/C/C37)



European Patent
Office

INCOMPLETE SEARCH
SHEET C

Application Number
EP 01 11 8978

Claim(s) searched completely:
17

Claim(s) searched incompletely:
1-16

Reason for the limitation of the search:

Present claims 1-16 relate to an extremely large number of possible compounds (any ester of ferulic acid). Support within the meaning of Article 84 EPC and/or disclosure within the meaning of Article 83 EPC is to be found, however, for only a very small proportion of the compounds claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts relating to the ferulic acid esters explicitly mentioned on page 7 of the description.



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number
EP 01 11 8978

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
X	CHENG J-T ET AL: "ANTIHYPERTENSIVE ACTIVITY OF PHENOLICS FROM THE FLOWER OF LONICERA JAPONICA" ZHONGHUA YAOXUE ZAZHI - CHINESE PHARMACEUTICAL JOURNAL, ZHONGGUO YAOXUEHUI, TAIPEI, TW, vol. 46, no. 6, 1994, pages 575-582, XP008007443 ISSN: 1016-1015	17	
Y	* table 1 * -----	1-16	
			TECHNICAL FIELDS SEARCHED (Int.Cl.7)

EPO FORM 1503 03.82 (P04C10)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 01 11 8978

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

26-03-2003

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 4707472	A	17-11-1987	JP 61040298 A	26-02-1986
			EP 0216927 A1	08-04-1987
			WO 8600312 A1	16-01-1986
WO 9801143	A	15-01-1998	DE 19627376 A1	08-01-1998
			AU 3541697 A	02-02-1998
			WO 9801143 A1	15-01-1998
			EP 0912189 A1	06-05-1999
			JP 2001504443 T	03-04-2001
			KR 2000022470 A	25-04-2000
			US 2002012708 A1	31-01-2002
US 5958417	A	28-09-1999	NONE	

EPC FORM P0159

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82